

What is claimed is:

1. A method of treating a disorder resulting from dopamine-related dysfunction, comprising the steps of:

administering to a patient a full D<sub>1</sub> agonist wherein said agonist has a  
5 half-life of less than 6 hours and wherein said agonist is administered at a dose resulting in a first plasma concentration of agonist capable of activating D<sub>1</sub> dopamine receptors to produce a therapeutic effect; and

reducing said agonist dose at least once every 24 hours to obtain a  
second lower plasma concentration of agonist wherein said second concentration of  
10 agonist results in suboptimal activation of D<sub>1</sub> dopamine receptors for a period of time sufficient to prevent induction of tolerance.

2. The method of claim 1 wherein the agonist is selected from the group consisting of dinapsoline, dinoxylone, dihydrexidine, other D<sub>1</sub> agonists, and analogs and derivatives of said agonists, and combinations thereof.

3. The method of claim 1 wherein the disorder is selected from the group consisting of Parkinson's disease, autism, attention deficit disorder, schizophrenia, restless leg syndrome, memory loss, and sexual dysfunction.

4. The method of claim 1 wherein said agonist is administered parenterally.

5. The method of claim 4 wherein said parenteral administration route is selected from the group consisting of intradermal, subcutaneous, intramuscular, intraperitoneal, intrathecal, and intravenous administration.

6. The method of claim 4 wherein said parenteral administration is achieved using a sustained or pulsatile or sustained release dosage form.

7. The method of claim 4 wherein said parenteral administration is achieved using a metering pump.

8. The method of claim 1 wherein said agonist is administered intranasally.

9. The method of claim 1 wherein said agonist is administered orally.

10. The method of claim 1 wherein said agonist is administered in combination with an antioxidant.

11. The method of claim 1 wherein the period of time for reducing said agonist dose to obtain said second plasma concentration of agonist is at least one hour per each 24-hour dosing period.

12. The method of claim 1 wherein the period of time for reducing said  
5 agonist dose to obtain said second plasma concentration of agonist is about one hour to about four hours per each 24-hour dosing period.